



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Statistical reports of States and cities of the United States—Yearly and monthly.

COLORADO—*Denver*.—Month of July, 1901. Estimated population, 150,000. Total number of deaths, 254, including diphtheria, 1; enteric fever, 1; measles, 4; scarlet fever, 4, and 62 from tuberculosis.

CONNECTICUT—*Bridgeport*.—Month of July, 1901. Estimated population, 71,000. Total number of deaths, 115, including diphtheria, 1; enteric fever, 1; scarlet fever, 1; and 14 from tuberculosis.

IOWA—*Davenport*.—Month of July, 1901. Census population, 35,254. Total number of deaths, 62, including 8 from tuberculosis.

MASSACHUSETTS—*Lawrence*.—Month of July, 1901. Census population, 62,559. Total number of deaths, 100, including diphtheria, 1; enteric fever, 1, and 10 from tuberculosis.

Worcester.—Month of July, 1901. Census population, 118,421. Total number of deaths, 196, including diphtheria, 2; enteric fever, 2, and 22 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended August 17, 1901, from 88 observers, indicate that cerebro-spinal meningitis, dysentery, remittent fever, and enteric fever were more prevalent, and inflammation of bowels, influenza, and whooping cough were less prevalent than in the preceding week. Cerebro-spinal meningitis was reported present at 7, whooping cough at 13, measles at 14, diphtheria at 17, smallpox at 42, scarlet fever at 46, enteric fever at 79, and phthisis pulmonalis at 195 places.

MINNESOTA—*St. Paul*.—Month of July, 1901. Estimated population, 165,000. Total number of deaths, 41, including diphtheria, 8; scarlet fever, 1, and 23 from tuberculosis.

NEW HAMPSHIRE—*Concord*.—Month of July, 1901. Estimated population, 19,500. Total number of deaths, 16. No deaths from contagious diseases.

NEW JERSEY—*Paterson*.—Month of July, 1901. Estimated population, 107,633. Total number of deaths, 196, including enteric fever, 3; whooping cough, 2, and 21 from phthisis pulmonalis.

NEW YORK—*Auburn*.—Month of July, 1901. Census population, 30,345. Total number of deaths, 33, including enteric fever, 1, and 4 from phthisis pulmonalis.

Rochester.—Month of July, 1901. Census population, 162,608. Total number of deaths, 186, including enteric fever, 2, and 10 from tuberculosis.

WASHINGTON—*Seattle*.—Month of July, 1901. Estimated population, 90,000. Total number of deaths, 69, including enteric fever, 1; smallpox, 2, and 12 from tuberculosis.